

Breast Milk Iodine and Perchlorate Concentrations in Lactating Boston-Area Women

Elizabeth N. Pearce, MD, MSc,¹ Angela M. Leung, MD,¹ Benjamin C. Blount, PhD,^{2*} Hamid R. Bazrafshan, MD,³ Xuemei He, MD,¹ Sam Pino,¹ Liza Valentin-Blasini, PhD,^{2*} Lewis E. Braverman, MD¹

¹ Section of Endocrinology, Diabetes, and Nutrition, Boston Medical Center and Boston University School of Medicine, Boston, MA

² Division of Laboratory Sciences, National Center for Environmental Health, Centers for Disease Control and Prevention, Atlanta, GA

³ Department of Medicine, Golestan University Medical School, Golestan, Islamic Republic of Iran

This is an un-copied author manuscript copyrighted by The Endocrine Society. This may not be duplicated or reproduced, other than for personal use or within the rule of “Fair Use of Copyrighted Materials” (section 107, Title 17, U.S. Code) without permission of the copyright owner, The Endocrine Society. From the time of acceptance following peer review, the full text of this manuscript is made freely available by The Endocrine Society at <http://www.endojournals.org/>. The final copy edited article can be found at <http://www.endojournals.org/>. The Endocrine Society disclaims any responsibility or liability for errors or omissions in this version of the manuscript or in any version derived from it by the National Institutes of Health or other parties. The citation of this article must include the following information: author(s), article title, journal title, year of publication and DOI.

*CDC’s role in this study dealt exclusively with perchlorate exposure assessment. The findings and conclusions in this report are those of the authors and do not necessarily represent the views of CDC.

Abbreviated title: Breast milk iodine and perchlorate

Key words: iodine, perchlorate, thyroid, lactation

Word count: 3,062

Number of figures: 3

Number of tables: 1

Corresponding Author and author for reprint requests: Elizabeth N. Pearce, M.D., M.Sc., Boston Medical Center, 88 East Newton Street, Evans 201, Boston, MA 02118; phone 617-414-1348, fax 617-638-7221, elizabeth.pearce@bmc.org

Abstract

Context: Breastfed infants rely on adequate maternal dietary iodine intake.

Objective: To measure breast milk iodine and perchlorate, an inhibitor of iodide transport into the thyroid and potentially into breast milk, in Boston-area women.

Participants: 57 lactating healthy volunteers in the Boston area.

Measurements: Breast milk iodine and perchlorate concentrations, and urine iodine, perchlorate, and cotinine concentrations were measured. For comparison, iodine and perchlorate levels in infant formulae were also measured.

Results: Median breast milk iodine content in 57 samples was 155 µg/L (range 2.7 – 1968 µg/L). Median urine iodine was 114 µg/L (range 25 – 920 µg/L). Perchlorate was detectable in all 49 breast milk samples (range 1.3 – 411 µg/L), all 56 urine samples (range 0.37 – 127 µg/L), and all 17 infant formulae samples (range 0.22 – 4.1 µg/L) measured. Breast milk iodine content was significantly correlated with urinary iodine per gram creatinine and urinary cotinine, but was not significantly correlated with breast milk or urinary perchlorate.

Conclusions: Perchlorate exposure was not significantly correlated with breast milk iodine concentrations. Perchlorate was detectable in infant formula, but at lower levels than in breast milk. 47% of women sampled may have been providing breast milk with insufficient iodine to meet infants' requirements.

Introduction

Thyroid hormone, requiring adequate iodine intake, is critical for neurodevelopment *in utero* and in early life. Worldwide, iodine deficiency remains the leading cause of preventable mental retardation (1). Since the 1920s, U.S. dietary iodine has generally been adequate. However, among U.S. women of childbearing age (15-44 years), median urinary iodine levels, a biomarker for dietary iodine, decreased by over 50% from 1971 to 1994 according to data from the National Health and Nutrition Examination Survey (NHANES) (2). More recent NHANES data indicate that urinary iodine levels have stabilized (3). Although the latest NHANES study demonstrated an adequate median urinary iodine level of 167.8 $\mu\text{g/L}$ in U.S. adults, 16.8% of U.S. women of childbearing age had urinary iodine concentrations $<50 \mu\text{g/L}$ (3). In our population at Boston Medical Center, we have reported that 9% of 100 women sampled had urinary iodine levels below $<50 \mu\text{g/L}$, and 49% had values below that recommended for pregnant women (4). Breastfed infants are reliant on adequate maternal dietary iodine intake (5). The Institute of Medicine's recommended dietary allowance for lactating women is 290 μg iodine daily (6). Based on concerns about adequate iodine intake in the perinatal period, the National Academy of Sciences recently recommended that consideration be given to adding iodine to all prenatal vitamins (7). The American Thyroid Association has also recently recommended that all women receive 150 μg iodine supplements daily during pregnancy and lactation and that all prenatal vitamins contain 150 μg of iodine (8).

Perchlorate is an anion that decreases the uptake of iodine into the thyroid gland and potentially into milk ducts in the lactating breast by competitively inhibiting the sodium/iodide symporter (NIS) on the basolateral surface of thyroid cells and on lactating breast cells.

Perchlorate is a component of solid rocket fuel, is found in Chilean nitrate fertilizers used in the

U.S., and is an environmental contaminant from natural processes in many regions of the U.S. Perchlorate has been detected in the drinking water of communities around the United States (9), including Massachusetts (10). It has also been detected in foods such as lettuce and wheat (11,12). Perchlorate exposure appears to be ubiquitous in the U.S. population (13). A recent study measured perchlorate in the breast milk of 36 women from 18 states and found detectable levels in all of the samples (range 0.6 - 92.2 $\mu\text{g/L}$) (14). Breast milk iodide and perchlorate concentrations were inversely correlated in the 6 samples with perchlorate concentrations ≥ 10 $\mu\text{g/L}$, although there were no correlations between breast milk iodide and perchlorate in the full data set (14).

Another recent study concluded that cigarette smoking decreases breast milk iodine concentrations (15). Cigarette smoke contains significant levels of cyanide that is metabolized to thiocyanate, a known competitive inhibitor of the NIS. Elevated serum thiocyanate levels may inhibit NIS-mediated transport of iodide in the lactating breast, leading to reduced breast milk iodine levels.

The objective of the present study was to determine whether breast milk iodine concentrations in Boston-area women are adequate for infant nutrition, and whether breast milk iodine concentrations may be associated with environmental perchlorate or cigarette smoke exposure. Levels of iodine and perchlorate in breast milk were compared with those measured in seventeen brands of infant formulae.

Materials and Methods

We obtained breast milk and urine samples from 57 healthy Boston-area volunteers (range, 19 – 45 years; mean age $30 \pm \text{SD } 6$ years), at 10 – 250 (median 48) days postpartum,

between July 2002 – April 2006. The local Institutional Review Board approved the protocol and informed consent was obtained from all participants. Volunteers were recruited through various means, including a community-based new mothers' group, routine post-partum visits at a hospital-based inner-city obstetric clinic, and through flyers posted in a hospital lobby. 72% women studied reported taking prenatal multivitamins, but only 3 were using iodine-containing multivitamin preparations.

27 women completed a questionnaire regarding their demographics and intake of dietary iodine and iodine-containing multivitamins. Spot urinary iodine measurements were obtained from all subjects. Urine perchlorate, creatinine, and cotinine (a metabolite of nicotine in cigarette smoke) concentrations were also measured. Samples of breast milk (approximately 10 mL) were collected at the start of a feed using either hand-expression or a breast pump in 27 of the women. Samples of breast milk were collected in 5 mL increments sequentially using a breast pump in the other 30 women. In this subset of 30 subjects, we measured breast milk iodine concentrations both at the start of a feed and sequentially throughout a single feed, in order to assess any potential intrafeed variation. In the women whose breast milk was collected in sequential increments, the mean breast milk iodine concentration is reported. Enough breast milk was available in 49 (86%) and enough urine was available in 56 (98%) of the samples for the measurement of perchlorate. Enough urine was available in 56 (98%) of the samples for the measurement of cotinine. All breast milk and urine samples were obtained within the same hour.

Seventeen brands of infant formulae were also assessed for iodine and perchlorate levels. A single sample of each different type of liquid formula available at a local supermarket was purchased for testing. Nine brands were sold in concentrated form and designed to be diluted by half before use. Iodine and perchlorate levels were measured directly in these samples, and the

results were divided by half to reflect the concentration intended for infant use. The other eight brands were sold ready for use.

Breast milk, infant formulae, and urine iodine concentrations were measured spectrophotometrically by a modification of the method of Benotti et al (16). Iodine concentrations were measured at least twice; in 95% of the samples, the initial two measurements were within 15% of each other and the two values were averaged. In the case where the initial two measurements were not within 15% of each other, a third measurement was obtained and the average of all measurements was reported. Cotinine measurements were performed by immunoassay (Immulite 2000 Nicotine Metabolite Assay; Diagnostic Products Corporation, Los Angeles, CA). Urine creatinine measurements were performed using Jaffe's alkaline picrate method. The perchlorate content of breast milk, infant formulae, and urine samples was measured at the Centers for Disease Control and Prevention laboratories in Atlanta, GA using ion chromatography-mass spectrometry (17).

All statistical analyses were carried out using SAS version 9.1 (SAS Institute, Cary, NC). Spearman's rank correlation coefficient was used to determine whether linear associations were present, and multivariate linear regression was used to determine significant predictors of breast milk iodine concentration. Intrafeed variation in the iodine content of the subjects whose breast milk was measured sequentially was assessed using repeated measures ANOVA. Differences in breast milk and urine iodine values between smokers and nonsmokers and differences in measured and labeled infant formula iodine content were assessed using an independent t test. Differences in mean values across subject groups were assessed by ANOVA. Differences in median iodine and perchlorate values in breast milk compared to infant formula were assessed using the Wilcoxon Rank Sum test.

Results

Median iodine content in the 57 breast milk samples was 155 $\mu\text{g/L}$ (range, 2.7 – 1968 $\mu\text{g/L}$; mean (\pm SD), 205 \pm 271 $\mu\text{g/L}$). There was no significant intrafeed variation in the iodine content of the subjects whose breast milk was assessed sequentially during a single feed. Breast milk iodine concentrations in the three women who reported using iodine-containing multivitamins were 28 $\mu\text{g/L}$, 68 $\mu\text{g/L}$, and 187 $\mu\text{g/L}$. In a subset of 27 women whose use of iodized table salt was determined, there was no significant difference in breast milk iodine levels between the subjects who reported regular, occasional, or no use of iodized salt ($p = 0.16$). There was a slight but significant positive correlation between breast milk iodine content and maternal age ($r^2 = 0.08$, $p = 0.04$), but not infant age ($p = 0.4$). Median urine iodine content was 114 $\mu\text{g/L}$ (range, 25 – 920 $\mu\text{g/L}$; mean, 155 \pm 142).

Perchlorate was detectable in all 49 breast milk samples (median, 9.1 µg/L; range, 1.3 – 411; mean, 33 ± 77) and 56 urine samples tested (median, 3.0 µg/L; range, 0.37 – 127; mean, 8.2 ± 19). Cotinine was detected in the urine of 32 (57%) of 56 urine samples tested (range 1.5 – 1575 ng/mL), with 6 (19%) of these likely to be smokers (urinary cotinine >500 ng/mL). Mean breast milk iodine was 62 ± 35 µg/L in the smokers and 221 ± 285 µg/L in the nonsmokers ($p = 0.0005$), while urinary iodine did not differ between these groups ($p = 1.0$). There were significant differences in the mean breast milk perchlorate concentrations between the women recruited from the mothers seen at routine post-partum visits at a hospital-based inner-city obstetric clinic ($n = 10$; 88.0 ± 150.0 µg/L), those from the community-based new mothers' group ($n = 9$; 50.0 ± 62.5 µg/L), and those recruited by the hospital postings ($n = 30$; 10.3 ± 13.8 µg/L) (p for difference across the 3 groups was 0.01). There were no significant differences between breast milk iodine, urinary iodine, urinary perchlorate, and urinary cotinine across these subject groups.

Breast milk and urine iodine content in µg/L were not significantly correlated ($r^2 = 0.06$, $p = 0.08$). A significant positive correlation between breast milk iodine and urine iodine content per gram creatinine was observed ($r^2 = 0.27$, $p < 0.0001$). There was a significant positive correlation between breast milk perchlorate and urine perchlorate concentrations ($r^2 = 0.11$, $p = 0.02$). There were no significant correlations between breast milk iodine and perchlorate concentrations ($n = 49$, $r^2 = 0.05$, $p = 0.1$) (Figure 1), including the 23 women whose breast milk perchlorate values were ≥ 10 µg/L ($r^2 = 0.002$, $p = 0.8$). Breast milk iodine and urine perchlorate concentrations ($r^2 = 0.004$, $p = 0.7$) were not significantly correlated. There was a slight but significant inverse correlation between breast milk iodine and urine cotinine concentrations ($r^2 = 0.13$, $p = 0.006$) (Figure 2). In multivariate models, breast milk perchlorate ($p = 0.9$), urine

perchlorate ($p = 0.4$), urine cotinine ($p = 0.2$), and baby age ($p = 0.6$) did not predict breast milk iodine levels. Assuming an average daily infant intake of 0.78 liters breast milk (1) and that these breast milk samples are representative, 27 of the 57 samples (47%) did not contain sufficient iodine to meet infants' Adequate Intake (AI) of 110-130 $\mu\text{g}/\text{day}$, as recommended by the Institute of Medicine (Figure 3).

In the seventeen brands of infant formulae, the median measured iodine concentration was 145 $\mu\text{g}/\text{L}$ (range 84 – 224 $\mu\text{g}/\text{L}$) and the median perchlorate concentration was 1.50 $\mu\text{g}/\text{L}$ (range 0.22 – 4.1 $\mu\text{g}/\text{L}$) (Table 1). Measured infant formula iodine concentrations were significantly higher than labeled amounts ($p < 0.0001$). There was no significant difference between the median breast milk and infant formulae iodine concentrations ($p = 1.0$), while the median breast milk perchlorate concentration was significantly higher than the median infant formula perchlorate concentration ($p < 0.0001$).

Discussion

Iodine is avidly concentrated in the lactating breast, due to increased expression of NIS during lactation (18). Breast milk iodine levels are highest in transitional milk (2-5 days postpartum), and decrease to stable levels by 10 days postpartum (18). We report that among the 30 subjects whose breast milk was sampled sequentially, there were no significant intrafeed variations in iodine concentration. A recent study suggests, however, that there is a substantial amount of day-to-day and diurnal variation in breast milk iodine excretion (19).

The median breast milk iodide level in a 1984 sample of 37 U.S. women was 178 $\mu\text{g}/\text{L}$ (20), which is similar to the median iodine value of 155 $\mu\text{g}/\text{L}$ observed in our study. However, the median breast milk iodine value in our study was far higher than the median values of breast

milk iodide (33.5 $\mu\text{g/L}$ and 55.2 $\mu\text{g/L}$) observed recently by Kirk et al in two samples of U.S. women (14,19). Some, but not all, of this difference may be due to our method of measuring total iodine in contrast to Kirk et al who measured only ionic iodine (iodide); however, approximately 80-90% of iodine in human milk is in the form of iodide (Sam Pino, unpublished data). The median breast milk iodine concentration observed in our study was also far higher than the mean value of 38.5 $\mu\text{g/L}$ reported in 55 Chilean women (21); the iodine content of breast milk from the Chilean women was measured by the same method as reported in the current manuscript (16). Tellez et al in the Chilean study (21) and Chan et al in an Australian study (22) reported that breast milk iodine levels correlate with urinary iodine per gram creatinine but not with unadjusted urinary iodine values, as seen in the present study.

Assuming breast milk intake of 0.78 L/day (6) as the only source of breastfed infant iodine nutrition, and that single samples are representative of daily breast milk iodine content, which may not be correct due to the day-to-day and diurnal variation in breast milk iodine content noted by Kirk et al (19), 47% of women sampled may have been providing breast milk with insufficient iodine to meet infants' daily requirements. The difference between median breast milk and measured infant formulae iodine levels was not significant ($p = 1.0$), however there was a significant difference between the labeled and measured mean infant formula iodine concentrations ($p < 0.0001$). Assuming an average infant formula intake of 0.86 L/day (for infants 2 months of age) (23), the labeled iodine concentrations in all 17 (100%) infant formula brands were insufficient to meet the Institute of Medicine's recommended AI of 110 $\mu\text{g/day}$, while 35% of the measured infant formula iodine concentrations contained insufficient iodine levels. Although clinically apparent iodine deficiency disorders are vanishingly rare in U.S. infants, it is possible that breast-fed infants whose mothers' milk is iodine deficient, as well as

some formula-fed infants, might experience subtle neurodevelopmental abnormalities. An upper limit for iodine tolerability has not been established in infants.

The question of whether low-level environmental perchlorate exposure has clinical consequences has been extremely controversial (24), and has been the topic of a recent National Academy of Sciences review (7). Median urine perchlorate was 3.6 $\mu\text{g/L}$ in 2,820 spot urine specimens in the most recent NHANES survey (2001-2002) (13), and there was a small positive correlation with serum TSH and a small inverse correlation with serum T4 values (25). The developing fetus and infant are likely to be most vulnerable to the adverse effects of perchlorate on thyroid function since they have the highest rate of thyroidal iodine turnover and they require adequate thyroid hormone for normal neurodevelopment. In this study, we found measurable perchlorate levels in 100% of 49 human milk samples tested, with a median value of 9.1 $\mu\text{g/L}$. This is higher than the levels reported by Kirk et al in two recent studies (median 3.3 $\mu\text{g/L}$ and 4.0 $\mu\text{g/L}$) (14,19) and in the infant formulae currently measured (median 1.50 $\mu\text{g/L}$). There was a significant difference in the perchlorate levels among the subjects when grouped by method of recruitment, which may be attributable to the relatively small sample size as well as the demographic differences of the study population. In data from NHANES 2001-2002, non-Hispanic Blacks had lower levels of urinary perchlorate than non-Hispanic whites (13). Perchlorate was also detectable in all 56 human urine samples tested, with median levels (2.90 $\mu\text{g/L}$, and 2.94 $\mu\text{g/g}$ creatinine) consistent with previously published data (17). The source of this perchlorate exposure is unknown and merits further investigation.

Kirk et al previously speculated that milk perchlorate $\geq 10 \mu\text{g/L}$ may lead to reduced milk iodide levels (14). Despite the fact that 47% of our samples contained perchlorate at concentrations $\geq 10 \mu\text{g/L}$, we found no correlations between milk perchlorate and milk iodine

content. Our results are consistent with the results of Lengemann who found dairy cattle milk iodine excretion unaffected by perchlorate doses < 0.1 mg/kg-day (26). A recent study of pregnant and lactating Chilean women consuming tap water known to contain naturally occurring perchlorate (~ 110 $\mu\text{g/L}$) also found no correlation between perchlorate exposure and breast milk iodine or newborn thyroid function tests (21). One possible reason for the lack of correlation in these studies might be the diurnal and day-to-day variation in breast milk concentrations of both iodine and perchlorate (19).

Thiocyanate, as a metabolite of cigarette smoking, is also a known inhibitor of the NIS. Laurberg et al reported that although both smokers and non-smokers had stable levels of urinary iodine post-partum, smokers (defined as those with urinary cotinine > 500 ng/mL) had lower concentrations of breast milk iodine and their infants had lower concentrations of urinary iodine (15). Similarly, we found a small but significant inverse correlation between breast milk iodine and urine cotinine concentrations and found that smokers had significantly lower breast milk iodine levels than nonsmokers. Breastfeeding women should clearly be counseled against smoking for this and many other reasons.

We believe that a larger study of breast milk iodine content and its relationship to perchlorate exposure and maternal iodine nutrition is warranted in U.S. women. Although in the present study we found no correlation between breast milk iodine and perchlorate concentrations, the relatively low levels of iodine and relatively high levels of perchlorate in human milk do raise concerns. One limitation of the study was that we do not know the impact of breast milk perchlorate on the thyroidal iodine uptake and thyroid function of the breastfed infants. The American Thyroid Association has recently identified as a research priority the measurement of breast milk iodine levels in U.S. women and correlation with maternal iodine nutrition and

factors such as smoking (27). Very few women in the present study reported routine use of an iodine-containing multivitamin. We feel that there should be more public awareness of the importance of dietary iodine in lactation, and that iodine (at least 150 μg) should be included in standard prenatal multivitamins consistent with the recent recommendations of the National Academy of Sciences (7) and the American Thyroid Association (8). It is important to note that epidemiologic studies consistently point to the value of human milk as the healthiest food for infants. Thus, the benefits of human milk outweigh possible effects of environmental toxicants present.

- ¹ **WHO, UNICEF and ICCIDD.** Assessment of the Iodine Deficiency Disorders and Monitoring their Elimination. A guide for programme managers. Geneva, 2001. WHO publication, WHO/NHD/01.1.
- ² **Hollowell JG, Staehling NW, Hannon WH, Flanders DW, Gunter EW, Maberly GF, Braverman LE, Pino S, Miller DT, Garbe PL, DeLozier DM, Jackson RJ** 1998 Iodine nutrition in the United States. Trends and public health implications: iodine excretion data from National Health and Nutrition Examination Surveys I and III (1971-1974 and 1988-1994). *J Clin Endocrinol Metab* 83: 3401-3408
- ³ **Caldwell KL, Jones RL, Hollowell JG** 2005 Urinary iodine concentration: United States National Health And Nutrition Examination Survey 2001-2002. *Thyroid* 15:692-699
- ⁴ **Pearce EN, Bazrafshan HR, He X, Pino S, Braverman LE** 2004 Dietary Iodine in Pregnant Women from the Boston, Massachusetts Area. *Thyroid* 14:327-8
- ⁵ **Semba RD, Delange F** 2001 Iodine in human milk: perspectives for infant health. *Nutr Rev* 59:269-278
- ⁶ **Food and Nutrition Board, Institute of Medicine** 2006 Dietary reference intakes. Washington, D.C.: National Academy Press; 320-7
- ⁷ **Committee to Assess the Health Implications of Perchlorate Ingestion, National Research Council** 2005 Health Implications of Perchlorate Ingestion. Washington, D.C.: National Academies Press.
- ⁸ **Becker DV, Braverman LE, Delange F, Dunn JT, Franklyn JA, Hollowell JG, Lamm SH, Mitchell ML, Pearce E, Robbins J, Rovet JF** 2006 Iodine supplementation for pregnancy and lactation- United States and Canada: recommendations of the American Thyroid Association. *Thyroid* 16:949-51

- ⁹ Unregulated Contaminant Monitoring Regulation (UCMR) data from public water systems; U.S. Environmental Protection Agency. Environmental Protection Agency, 2004. <http://www.epa.gov/safewater/standard/ucmr/main.html>. (accessed 9/5/2006)
- ¹⁰ Draft Report: The Occurrence of Sources of Perchlorate in Massachusetts. Massachusetts Department of Environmental Protection, 2005. www.mass.gov/dep/cleanup/sites/percsour.pdf. (accessed 9/5/2006)
- ¹¹ **Yu L, Canas JE, Cobb GP, Jackson WA, Anderson TA** 2004 Uptake of perchlorate in terrestrial plants. *Ecotoxicol Environ Saf* 58:44-49
- ¹² **Jackson WA, Joseph P, Laxman P, Tan K, Smith PN, Yu L, Anderson TA** 2005 Perchlorate accumulation in forage and edible vegetation. *J Agric Food Chem* 53:369-373
- ¹³ **Blount BC, Valentin-Blasini L, Osterloh JD, Mauldin JP, Pirkle JL** 2006 Perchlorate Exposure of the US Population, 2001-2002. *J Exposure Sci Environ Epidemiol*. Doi: 10.1038/sj.jes.7500535
- ¹⁴ **Kirk AB, Martinelango PK, Tian K, Dutta A, Smith EE, Dasgupta PK** 2005 Perchlorate and iodide in dairy and breast milk. *Environ Sci Technol* 39:2011-2017
- ¹⁵ **Laurberg P, Nohr SB, Pedersen KM, Fuglsang E** 2004 Iodine nutrition in breast-fed infants is impaired by maternal smoking. *J Clin Endocrinol Metab* 89:181-187
- ¹⁶ **Benotti J, Benotti N, Pino S, Gardyna H** 1965 Determination of total iodine in urine, stool, diets, and tissue. *Clin Chem* 11:932-936
- ¹⁷ **Valentin-Blasini L, Mauldin JP, Maple D, Blount BC** 2005 Analysis of perchlorate in human urine using ion chromatography and electrospray tandem mass spectrometry. *Anal Chem* 77:2475-2481

-
- ¹⁸ **Etling N, Padovani E, Fouque F, Tato L** 1986 First-month variations in total iodine content of human breast milks. *Early Hum Dev* 13:81-85
- ¹⁹ **Kirk AB, Dyke JV, Martin CF, Dasgupta PK** 2006 Temporal Patterns in Perchlorate, Thiocyanate and Iodide Excretion in Human Milk. *Environmental Health Perspectives*.
Doi:10.1289/ehp.9558
- ²⁰ **Gushurst CA, Mueller JA, Green JA, Sedor F** 1984 Breast milk iodine reassessment in the 19890s. *Pediatrics* 73: 354-357
- ²¹ **Tellez R, Michaud Chacon P, Reyes Abarca C, Blount BC, Van Landingham CB, Crump KS, Gibbs JP** 2005 Long-term environmental exposure to perchlorate in drinking water and thyroid function during pregnancy and the neonatal period. *Thyroid* 15:963-975
- ²² **Chan SS, Hams G, Wiley V, Wilcken B, McElduff A** 2003 Postpartum maternal iodine status and the relationship to neonatal thyroid function. *Thyroid* 13:873-876
- ²³ **Anderson WA, Pratt I, Ryan MR, Flynn A** 2004 A Probabilistic estimate of fluoride intake by infants up to the age of 4 months from infant formula reconstituted with tap water in the fluorinated regions of Ireland. *Caries Research* 38:421-9
- ²⁴ **Hershman JM** 2005 Perchlorate and thyroid function: what are the environmental issues? *Thyroid* 15:427-431
- ²⁵ **Blount BC, Pirkle JL, Osterloh JD, Valentin-Blasini L, Caldwell KL** 2006 Urinary Perchlorate and Thyroid Hormone Levels in Adolescent and Adult Men and Women Living in the United States. *Environmental Health Perspectives* 114:1865-1871
- ²⁶ **Lengemann FW** 1973 Reduction of iodine transfer to milk of cows after perchlorate ingestion. *J Dairy Science* 56:753-756

²⁷ **American Thyroid Association** 2005 Consensus Statement #2: American Thyroid Association statement on early maternal thyroidal insufficiency: recognition, clinical management and research directions. *Thyroid* 15:77-79

Legends:

Figure 1. Correlation between breast milk iodine and perchlorate concentrations.

Figure 2. Correlation between breast milk iodine and urine cotinine concentrations.

Figure 3. Mean breast milk iodine content by quartile, with standard deviation (SD) shown numerically. Dotted line corresponds to approximate breast milk iodine content required to achieve Adequate Intake (110 – 130 $\mu\text{g/L}$) for an infant age 0-6 months.

Table 1:

Liquid infant formula brand	Labeled Iodine (µg/L)	Measured Iodine (µg/L)	Measured Perchlorate µg/L)
Milk-based			
Enfamil A.R. Lipil (milk-based, thickened with rice starch, iron-fortified)	67	224	4.1
Enfamil LactoFree Lipil (milk-based, lactose-free, iron-fortified) *	100	120	0.4
Enfamil Lipil Low Iron (milk-based)	67	158	2.1
Enfamil Lipil with Iron (milk-based infant formula) *	67	102	1.7
Enfamil with Iron (milk-based) *	67	162	2.0
Nestle Good Start with Iron (milk-based, Supreme, with easy to digest Comfort Proteins) *	80	145	0.3
Nestle Good Start with Iron (milk-based, Supreme, with easy to digest Comfort Proteins)	80	178	0.2
Nutramigen Lipil (milk-based, lactose-free, hypoallergenic, iron-fortified)	100	193	1.5
Similac Advanced Infant Formula with Iron (milk-based) *	40	122	1.5
Similac Alimentum Advance with Iron (milk-based, protein hydrolysate formula with iron, hypoallergenic)	100	158	2.0
Similac Infant Formula Low Iron (milk-based)	40	153	2.1

Similac Infant Formula with Iron (milk-based) *	40	142	1.6
Similac Lactose Free Infant Formula with Iron (milk-based)	60	143	1.4
Similac Neosure Advance with Iron (milk-based)	100	178	2.5
Soy-based			
Enfamil Prosobee Lipil (milk-free, lactose-free, iron-fortified) *	100	112	0.3
Enfamil Prosobee Soy Infant Formula (milk-free, lactose-free, iron-fortified) *	100	122	0.6
Similac Isomil Soy Formula with Iron (milk-free, lactose-free)*	100	84	0.4

Table 1. Iodine and perchlorate concentrations in seventeen brands of infant formulae. The nine brands of infant formula that were sold in concentrated form and designed to be diluted by half before use are denoted by the *. All of the iodine and perchlorate levels reported here reflect the concentration intended for infant use.

Figure 1:




Figure 2:




Figure 3:

